

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 November 2001 (08.11.2001)

PCT

(10) International Publication Number
WO 01/83745 A1

(51) International Patent Classification⁷: **C12N 15/12**,
C07K 14/47, C12N 15/62, C07K 16/18, G01N 33/68,
C12Q 1/68

2a, 63477 Maintal (DE). **HOHEISEL, Jörg, Dietrich**
[DE/DE]; Lempenseite 18-5, 69168 Wiesloch (DE).
FROHME, Marcus [DE/DE]; Danzingerstrasse 6, 68535
Edingen (DE).

(21) International Application Number: PCT/EP01/04888

(74) Common Representative: **MERCK PATENT GMBH**;
Frankfurter Strasse 250, 64293 Darmstadt (DE).

(22) International Filing Date: 30 April 2001 (30.04.2001)

(25) Filing Language:

English

(81) Designated States (*national*): CA, JP, US.

(26) Publication Language:

English

(84) Designated States (*regional*): European patent (AT, BE,
CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC,
NL, PT, SE, TR).

(30) Priority Data:

00109079.4

2 May 2000 (02.05.2000) EP

Published:

— with international search report


— before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant (*for all designated States except US*): **MERCK
PATENT GMBH** [DE/DE]; Frankfurter Strasse 250,
64293 Darmstadt (DE).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **SIRRENBURG,**
Christian [DE/DE]; Heinrich-Delp-Strasse 221, 64297
Darmstadt (DE). **HOCK, Björn** [DE/DE]; Nordstrasse

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*



WO 01/83745 A1

(54) Title: NEW GTPASE-ACTIVATING PROTEIN 1

(57) Abstract: New GTPase-Activating Protein 1 polypeptides and polynucleotides and methods for producing such polypeptides by recombinant techniques are disclosed. Also disclosed are methods for utilizing New GTPase-Activating Protein 1 polypeptides and polynucleotides in diagnostic assays.

New GTPase-activating protein 1

Field of the Invention

This invention relates to newly identified polypeptides and polynucleotides encoding such polypeptides sometimes hereinafter referred to as „New GTPase-Activating Protein 1 (NGAP-1)“, to their use in diagnosis and in identifying compounds that may be agonists, antagonists that are potentially useful in therapy, and to production of such polypeptides and polynucleotides.

Background of the Invention

The drug discovery process is currently undergoing a fundamental revolution as it embraces "functional genomics", that is, high throughput genome- or gene-based biology. This approach as a means to identify genes and gene products as therapeutic targets is rapidly superceding earlier approaches based on "positional cloning". A phenotype, that is a biological function or genetic disease, would be identified and this would then be tracked back to the responsible gene, based on its genetic map position.

Functional genomics relies heavily on high-throughput DNA sequencing technologies and the various tools of bioinformatics to identify gene sequences of potential interest from the many molecular biology databases now available. There is a continuing need to identify and characterise further genes and their related polypeptides/proteins, as targets for drug discovery.

Members of the RAS superfamily of GTPases are e.g. involved in growth control. Certain GTP-binding protein mutants e.g. v-ras are oncoproteins. GTPase activating proteins enhance the intrinsically slow GTPase activity of Ras-like proteins, which function as molecular switches controlling cellular pathways. GAPs turn off these switches by stimulating the GTPase activity of Ras-like proteins. Loss of GAP function is e.g. responsible for the disease phenotype in type 1 neurofibromatosis patients.

The amino acid sequence of the New GTPase-Activating Protein 1 is related to spindle checkpoint proteins, e.g. BUB2p and Cdc16, and to the *Drosophila* adhesion molecule pollux (Neuwald, A.F. (1997), *Trends Biochem. Sci.* **22**, 243-244).

5

Summary of the Invention

The present invention relates to New GTPase-Activating Protein 1, in particular New GTPase-Activating Protein 1 polypeptides and New GTPase-Activating Protein 1 polynucleotides, recombinant materials and methods for their production. Such polypeptides and polynucleotides are of interest in relation to methods of treatment of certain diseases, including, but not limited to, cancer, neuronal diseases, neurofibromatosis, hereinafter referred to as "diseases of the invention". In a further aspect, the invention relates to methods for identifying agonists and antagonists (e.g., inhibitors) using the materials provided by the invention, and treating conditions associated with New GTPase-Activating Protein 1 imbalance with the identified compounds. In a still further aspect, the invention relates to diagnostic assays for detecting diseases associated with inappropriate New GTPase-Activating Protein 1 activity or levels.

20

Description of the Invention

In a first aspect, the present invention relates to New GTPase-Activating Protein 1 polypeptides. Such polypeptides include:

(a) a polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1;

25

(b) a polypeptide comprising a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;

(c) a polypeptide comprising the polypeptide sequence of SEQ ID NO:2;

30

(d) a polypeptide having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;

- 3 -

(e) the polypeptide sequence of SEQ ID NO:2; and

(f) a polypeptide having or comprising a polypeptide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO:2;

5 (g) fragments and variants of such polypeptides in (a) to (f).

Polypeptides of the present invention are believed to be members of the GAP family and therefore may suppress the formation of tumors. Furthermore New GTPase-Activating Protein 1 may be part of a cell cycle checkpoint control system.

10 The biological properties of the New GTPase-Activating Protein 1 are hereinafter referred to as "biological activity of New GTPase-Activating Protein 1" or "New GTPase-Activating Protein 1 activity". Preferably, a polypeptide of the present invention exhibits at least one biological activity of New GTPase-Activating Protein 1.

15 Polypeptides of the present invention also includes variants of the aforementioned polypeptides, including all allelic forms and splice variants. Such polypeptides vary from the reference polypeptide by insertions, deletions, and substitutions that may be conservative or non-conservative, or any combination thereof. Particularly preferred variants are those in
20 which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acids are inserted, substituted, or deleted, in any combination.

Preferred fragments of polypeptides of the present invention include a polypeptide comprising an amino acid sequence having at least 30, 50 or
25 100 contiguous amino acids from the amino acid sequence of SEQ ID NO: 2, or a polypeptide comprising an amino acid sequence having at least 30, 50 or 100 contiguous amino acids truncated or deleted from the amino acid sequence of SEQ ID NO: 2. Preferred fragments are biologically active fragments that mediate the biological activity of New
30 GTPase-Activating Protein 1, including those with a similar activity or an improved activity, or with a decreased undesirable activity. Also preferred are those fragments that are antigenic or immunogenic in an animal, especially in a human.

- 4 -

Fragments of the polypeptides of the invention may be employed for producing the corresponding full-length polypeptide by peptide synthesis; therefore, these variants may be employed as intermediates for producing the full-length polypeptides of the invention. The polypeptides of the present invention may be in the form of the "mature" protein or may be a part of a larger protein such as a precursor or a fusion protein. It is often advantageous to include an additional amino acid sequence that contains secretory or leader sequences, pro-sequences, sequences that aid in purification, for instance multiple histidine residues, or an additional sequence for stability during recombinant production.

Polypeptides of the present invention can be prepared in any suitable manner, for instance by isolation from naturally occurring sources, from genetically engineered host cells comprising expression systems (*vide infra*) or by chemical synthesis, using for instance automated peptide synthesizers, or a combination of such methods. Means for preparing such polypeptides are well understood in the art.

In a further aspect, the present invention relates to New GTPase-Activating Protein 1 polynucleotides. Such polynucleotides include:

- (a) a polynucleotide comprising a polynucleotide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide sequence of SEQ ID NO:1;
- (b) a polynucleotide comprising the polynucleotide of SEQ ID NO:1;
- (c) a polynucleotide having at least 95%, 96%, 97%, 98%, or 99% identity to the polynucleotide of SEQ ID NO:1;
- (d) the polynucleotide of SEQ ID NO:1;
- (e) a polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;
- (f) a polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2;

- 5 -

(g) a polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95%, 96%, 97%, 98%, or 99% identity to the polypeptide sequence of SEQ ID NO:2;

(h) a polynucleotide encoding the polypeptide of SEQ ID NO:2;

5 (i) a polynucleotide having or comprising a polynucleotide sequence that has an Identity Index of 0.95, 0.96, 0.97, 0.98, or 0.99 compared to the polynucleotide sequence of SEQ ID NO:1;

(j) a polynucleotide having or comprising a polynucleotide sequence encoding a polypeptide sequence that has an Identity Index of 0.95, 0.96,
10 0.97, 0.98, or 0.99 compared to the polypeptide sequence of SEQ ID NO:2; and

polynucleotides that are fragments and variants of the above mentioned polynucleotides or that are complementary to above mentioned polynucleotides, over the entire length thereof.

15 Preferred fragments of polynucleotides of the present invention include a polynucleotide comprising an nucleotide sequence having at least 15, 30, 50 or 100 contiguous nucleotides from the sequence of SEQ ID NO: 1, or a polynucleotide comprising an sequence having at least 30, 50 or 100 contiguous nucleotides truncated or deleted from the sequence of SEQ
20 ID NO: 1.

Preferred variants of polynucleotides of the present invention include splice variants, allelic variants, and polymorphisms, including polynucleotides having one or more single nucleotide polymorphisms (SNPs).

25 Polynucleotides of the present invention also include polynucleotides encoding polypeptide variants that comprise the amino acid sequence of SEQ ID NO:2 and in which several, for instance from 50 to 30, from 30 to 20, from 20 to 10, from 10 to 5, from 5 to 3, from 3 to 2, from 2 to 1 or 1 amino acid residues are substituted, deleted or added, in any combination.

30 In a further aspect, the present invention provides polynucleotides that are RNA transcripts of the DNA sequences of the present invention. Accordingly, there is provided an RNA polynucleotide that:

- 6 -

(a) comprises an RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO:2;

(b) is the RNA transcript of the DNA sequence encoding the polypeptide of SEQ ID NO:2;

5 (c) comprises an RNA transcript of the DNA sequence of SEQ ID NO:1; or

(d) is the RNA transcript of the DNA sequence of SEQ ID NO:1;

and RNA polynucleotides that are complementary thereto.

10 The polynucleotide sequence of SEQ ID NO:1 shows homology with tre-2 oncogene (Acc.: S22155); spindle assembly checkpoint protein BUB2 (Acc.: P26448); yeast cell cycle regulators cdc16 (Acc.: P36618); human Rab6 GAP and centrosome-associated (Acc.: AJ011679); yeast Rab GAP Gyp1 (Acc.: S66953); Drosophila pullux protein (Acc.: Y17919); M. The
15 polynucleotide sequence of SEQ ID NO:1 is a cDNA sequence that encodes the polypeptide of SEQ ID NO:2. The polynucleotide sequence encoding the polypeptide of SEQ ID NO:2 may be identical to the polypeptide encoding sequence of SEQ ID NO:1 or it may be a sequence other than SEQ ID NO:1, which, as a result of the redundancy
20 (degeneracy) of the genetic code, also encodes the polypeptide of SEQ ID NO:2. The polypeptide of the SEQ ID NO:2 is related to other proteins of the GTPase-Activating Protein family, having homology and/or structural similarity with tre-2 oncogene (Acc.: S22155); spindle assembly checkpoint protein BUB2 (Acc.: P26448); yeast cell cycle regulators cdc16 (Acc.:
25 P36618); Drosophila pullux protein (Acc.:); M. musculus Tbc1 (Acc.:g988221).

Preferred polypeptides and polynucleotides of the present invention are expected to have, *inter alia*, similar biological functions/properties to their homologous polypeptides and polynucleotides. Furthermore, preferred
30 polypeptides and polynucleotides of the present invention have at least one New GTPase-Activating Protein 1 activity.

Polynucleotides of the present invention may be obtained using standard cloning and screening techniques from a cDNA library derived from mRNA in cells of human palate, tongue, testis, lung, neutrophils, activated macrophages, pancreas tumor, colon tumor, (see for instance, Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (1989)). Polynucleotides of the invention can also be obtained from natural sources such as genomic DNA libraries or can be synthesized using well known and commercially available techniques.

When polynucleotides of the present invention are used for the recombinant production of polypeptides of the present invention, the polynucleotide may include the coding sequence for the mature polypeptide, by itself, or the coding sequence for the mature polypeptide in reading frame with other coding sequences, such as those encoding a leader or secretory sequence, a pre-, or pro- or prepro- protein sequence, or other fusion peptide portions. For example, a marker sequence that facilitates purification of the fused polypeptide can be encoded. In certain preferred embodiments of this aspect of the invention, the marker sequence is a hexa-histidine peptide, as provided in the pQE vector (Qiagen, Inc.) and described in Gentz *et al.*, Proc Natl Acad Sci USA (1989) 86:821-824, or is an HA tag. The polynucleotide may also contain non-coding 5' and 3' sequences, such as transcribed, non-translated sequences, splicing and polyadenylation signals, ribosome binding sites and sequences that stabilize mRNA.

Polynucleotides that are identical, or have sufficient identity to a polynucleotide sequence of SEQ ID NO:1, may be used as hybridization probes for cDNA and genomic DNA or as primers for a nucleic acid amplification reaction (for instance, PCR). Such probes and primers may be used to isolate full-length cDNAs and genomic clones encoding polypeptides of the present invention and to isolate cDNA and genomic clones of other genes (including genes encoding paralogs from human sources and orthologs and paralogs from species other than human) that have a high sequence similarity to SEQ ID NO:1, typically at least 95% identity. Preferred probes and primers will generally comprise at least 15 nucleotides, preferably, at least 30 nucleotides and may have at least 50, if not at least 100 nucleotides. Particularly preferred probes will have

between 30 and 50 nucleotides. Particularly preferred primers will have between 20 and 25 nucleotides.

5 A polynucleotide encoding a polypeptide of the present invention, including homologs from species other than human, may be obtained by a process comprising the steps of screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof, preferably of at least 15 nucleotides; and isolating full-length cDNA and genomic clones containing said polynucleotide sequence. Such hybridization techniques are well known to the skilled artisan. Preferred stringent hybridization conditions include overnight incubation at 10 42°C in a solution comprising: 50% formamide, 5xSSC (150mM NaCl, 15mM trisodium citrate), 50 mM sodium phosphate (pH7.6), 5x Denhardt's solution, 10 % dextran sulfate, and 20 microgram/ml denatured, sheared salmon sperm DNA; followed by washing the filters in 0.1x SSC at about 15 65°C. Thus the present invention also includes polynucleotides, preferably with a nucleotide sequence of at least 100, obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO:1 or a fragment thereof, preferably of at least 15 nucleotides.

20 The skilled artisan will appreciate that, in many cases, an isolated cDNA sequence will be incomplete, in that the region coding for the polypeptide does not extend all the way through to the 5' terminus. This is a consequence of reverse transcriptase, an enzyme with inherently low "processivity" (a measure of the ability of the enzyme to remain attached 25 to the template during the polymerisation reaction), failing to complete a DNA copy of the mRNA template during first strand cDNA synthesis.

There are several methods available and well known to those skilled in the art to obtain full-length cDNAs, or extend short cDNAs, for example those based on the method of Rapid Amplification of cDNA ends (RACE) 30 (see, for example, Frohman et al., Proc Nat Acad Sci USA 85, 8998-9002, 1988). Recent modifications of the technique, exemplified by the Marathon (trademark) technology (Clontech Laboratories Inc.) for example, have significantly simplified the search for longer cDNAs. In the Marathon (trade mark) technology, cDNAs have been prepared from mRNA extracted from a chosen tissue and an 'adaptor' sequence ligated 35 onto each end. Nucleic acid amplification (PCR) is then carried out to

amplify the "missing" 5' end of the cDNA using a combination of gene specific and adaptor specific oligonucleotide primers. The PCR reaction is then repeated using 'nested' primers, that is, primers designed to anneal within the amplified product (typically an adaptor specific primer that anneals further 3' in the adaptor sequence and a gene specific primer that anneals further 5' in the known gene sequence). The products of this reaction can then be analysed by DNA sequencing and a full-length cDNA constructed either by joining the product directly to the existing cDNA to give a complete sequence, or carrying out a separate full-length PCR using the new sequence information for the design of the 5' primer.

Recombinant polypeptides of the present invention may be prepared by processes well known in the art from genetically engineered host cells comprising expression systems. Accordingly, in a further aspect, the present invention relates to expression systems comprising a polynucleotide or polynucleotides of the present invention, to host cells which are genetically engineered with such expression systems and to the production of polypeptides of the invention by recombinant techniques. Cell-free translation systems can also be employed to produce such proteins using RNAs derived from the DNA constructs of the present invention.

For recombinant production, host cells can be genetically engineered to incorporate expression systems or portions thereof for polynucleotides of the present invention. Polynucleotides may be introduced into host cells by methods described in many standard laboratory manuals, such as Davis et al., Basic Methods in Molecular Biology (1986) and Sambrook *et al.* (*ibid*). Preferred methods of introducing polynucleotides into host cells include, for instance, calcium phosphate transfection, DEAE-dextran mediated transfection, transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, scrape loading, ballistic introduction or infection.

Representative examples of appropriate hosts include bacterial cells, such as *Streptococci*, *Staphylococci*, *E. coli*, *Streptomyces* and *Bacillus subtilis* cells; fungal cells, such as yeast cells and *Aspergillus* cells; insect cells

- 10 -

such as *Drosophila* S2 and *Spodoptera* Sf9 cells; animal cells such as CHO, COS, HeLa, C127, 3T3, BHK, HEK 293 and Bowes melanoma cells; and plant cells.

5 A great variety of expression systems can be used, for instance, chromosomal, episomal and virus-derived systems, e.g., vectors derived from bacterial plasmids, from bacteriophage, from transposons, from yeast episomes, from insertion elements, from yeast chromosomal elements, from viruses such as baculoviruses, papova viruses, such as SV40, vaccinia viruses, adenoviruses, fowl pox viruses, pseudorabies viruses and
10 retroviruses, and vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, such as cosmids and phagemids. The expression systems may contain control regions that regulate as well as engender expression. Generally, any system or vector that is able to maintain, propagate or express a polynucleotide to produce a polypeptide in a host may be used. The appropriate polynucleotide sequence may be inserted into an expression system by any of a variety of well-known and routine techniques, such as, for example, those set forth in Sambrook *et al.*, (*ibid*). Appropriate secretion
15 signals may be incorporated into the desired polypeptide to allow secretion of the translated protein into the lumen of the endoplasmic reticulum, the periplasmic space or the extracellular environment. These signals may be endogenous to the polypeptide or they may be heterologous signals.

If a polypeptide of the present invention is to be expressed for use in screening assays, it is generally preferred that the polypeptide be
25 produced at the surface of the cell. In this event, the cells may be harvested prior to use in the screening assay. If the polypeptide is secreted into the medium, the medium can be recovered in order to recover and purify the polypeptide. If produced intracellularly, the cells must first be lysed before the polypeptide is recovered.

30 Polypeptides of the present invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxylapatite
35 chromatography and lectin chromatography. Most preferably, high performance liquid chromatography is employed for purification. Well

- 11 -

known techniques for refolding proteins may be employed to regenerate active conformation when the polypeptide is denatured during intracellular synthesis, isolation and/or purification.

5 Polynucleotides of the present invention may be used as diagnostic reagents, through detecting mutations in the associated gene. Detection of a mutated form of the gene characterised by the polynucleotide of SEQ ID NO:1 in the cDNA or genomic sequence and which is associated with a dysfunction will provide a diagnostic tool that can add to, or define, a diagnosis of a disease, or susceptibility to a disease, which results from
10 under-expression, over-expression or altered spatial or temporal expression of the gene. Individuals carrying mutations in the gene may be detected at the DNA level by a variety of techniques well known in the art.

15 Nucleic acids for diagnosis may be obtained from a subject's cells, such as from blood, urine, saliva, tissue biopsy or autopsy material. The genomic DNA may be used directly for detection or it may be amplified enzymatically by using PCR, preferably RT-PCR, or other amplification techniques prior to analysis. RNA or cDNA may also be used in similar fashion. Deletions and insertions can be detected by a change in size of the amplified product in comparison to the normal genotype. Point mutations can be identified by
20 hybridizing amplified DNA to labeled New GTPase-Activating Protein 1 nucleotide sequences. Perfectly matched sequences can be distinguished from mismatched duplexes by RNase digestion or by differences in melting temperatures. DNA sequence difference may also be detected by alterations in the electrophoretic mobility of DNA fragments in gels, with or
25 without denaturing agents, or by direct DNA sequencing (see, for instance, Myers *et al.*, Science (1985) 230:1242). Sequence changes at specific locations may also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (see Cotton *et al.*, Proc Natl Acad Sci USA (1985) 85: 4397-4401).

30 An array of oligonucleotide probes comprising New GTPase-Activating Protein 1 polynucleotide sequence or fragments thereof can be constructed to conduct efficient screening of *e.g.*, genetic mutations. Such arrays are preferably high density arrays or grids. Array technology methods are well known and have general applicability and can be used to
35 address a variety of questions in molecular genetics including gene expression, genetic linkage, and genetic variability, see, for example,

M.Chee et al., Science, 274, 610-613 (1996) and other references cited therein.

Detection of abnormally decreased or increased levels of polypeptide or mRNA expression may also be used for diagnosing or determining susceptibility of a subject to a disease of the invention. Decreased or increased expression can be measured at the RNA level using any of the methods well known in the art for the quantitation of polynucleotides, such as, for example, nucleic acid amplification, for instance PCR, RT-PCR, RNase protection, Northern blotting and other hybridization methods. Assay techniques that can be used to determine levels of a protein, such as a polypeptide of the present invention, in a sample derived from a host are well-known to those of skill in the art. Such assay methods include radioimmunoassays, competitive-binding assays, Western Blot analysis and ELISA assays.

Thus in another aspect, the present invention relates to a diagnostic kit comprising:

(a) a polynucleotide of the present invention, preferably the nucleotide sequence of SEQ ID NO: 1, or a fragment or an RNA transcript thereof;

(b) a nucleotide sequence complementary to that of (a);

(c) a polypeptide of the present invention, preferably the polypeptide of SEQ ID NO:2 or a fragment thereof; or

(d) an antibody to a polypeptide of the present invention, preferably to the polypeptide of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component. Such a kit will be of use in diagnosing a disease or susceptibility to a disease, particularly diseases of the invention, amongst others.

The polynucleotide sequences of the present invention are valuable for chromosome localisation studies. The sequence is specifically targeted to, and can hybridize with, a particular location on an individual human chromosome. The mapping of relevant sequences to chromosomes

according to the present invention is an important first step in correlating those sequences with gene associated disease. Once a sequence has been mapped to a precise chromosomal location, the physical position of the sequence on the chromosome can be correlated with genetic map data. Such data are found in, for example, V. McKusick, Mendelian Inheritance in Man (available on-line through Johns Hopkins University Welch Medical Library). The relationship between genes and diseases that have been mapped to the same chromosomal region are then identified through linkage analysis (co-inheritance of physically adjacent genes). Precise human chromosomal localisations for a genomic sequence (gene fragment etc.) can be determined using Radiation Hybrid (RH) Mapping (Walter, M. Spillelt, D., Thomas, P., Weissenbach, J., and Goodfellow, P., (1994) A method for constructing radiation hybrid maps of whole genomes, *Nature Genetics* 7, 22-28). A number of RH panels are available from Research Genetics (Huntsville, AL, USA) e.g. the GeneBridge4 RH panel (Hum Mol Genet 1996 Mar;5(3):339-46 A radiation hybrid map of the human genome. Gyapay G, Schmitt K, Fizames C, Jones H, Vega-Czarny N, Spillelt D, Muselet D, Prud'Homme JF, Dib C, Auffray C, Morissette J, Weissenbach J, Goodfellow PN). To determine the chromosomal location of a gene using this panel, 93 PCRs are performed using primers designed from the gene of interest on RH DNAs. Each of these DNAs contains random human genomic fragments maintained in a hamster background (human / hamster hybrid cell lines). These PCRs result in 93 scores indicating the presence or absence of the PCR product of the gene of interest. These scores are compared with scores created using PCR products from genomic sequences of known location. This comparison is conducted at <http://www.genome.wi.mit.edu/>. The gene of the present invention maps to human chromosome 9.

The polynucleotide sequences of the present invention are also valuable tools for tissue expression studies. Such studies allow the determination of expression patterns of polynucleotides of the present invention which may give an indication as to the expression patterns of the encoded polypeptides in tissues, by detecting the mRNAs that encode them. The techniques used are well known in the art and include in situ hybridisation techniques to clones arrayed on a grid, such as cDNA microarray

- 14 -

hybridisation (Schena *et al*, Science, 270, 467-470, 1995 and Shalon *et al*,
Genome Res; 6, 639-645, 1996) and nucleotide amplification techniques
such as PCR. A preferred method uses the TAQMAN (Trade mark)
technology available from Perkin Elmer. Results from these studies can
5 provide an indication of the normal function of the polypeptide in the
organism. In addition, comparative studies of the normal expression
pattern of mRNAs with that of mRNAs encoded by an alternative form of
the same gene (for example, one having an alteration in polypeptide coding
potential or a regulatory mutation) can provide valuable insights into the role
10 of the polypeptides of the present invention, or that of inappropriate
expression thereof in disease. Such inappropriate expression may be of a
temporal, spatial or simply quantitative nature.

The polypeptides of the present invention are expressed in testis, tongue,
palate, neutrophils, activated monocytes, pancreas, tumor, colon tumor.

15

A further aspect of the present invention relates to antibodies. The
polypeptides of the invention or their fragments, or cells expressing them,
can be used as immunogens to produce antibodies that are immunospecific
for polypeptides of the present invention. The term "immunospecific"
20 means that the antibodies have substantially greater affinity for the
polypeptides of the invention than their affinity for other related polypeptides
in the prior art.

Antibodies generated against polypeptides of the present invention may be
obtained by administering the polypeptides or epitope-bearing fragments, or
cells to an animal, preferably a non-human animal, using routine protocols.
25 For preparation of monoclonal antibodies, any technique which provides
antibodies produced by continuous cell line cultures can be used.
Examples include the hybridoma technique (Kohler, G. and Milstein, C.,
Nature (1975) 256:495-497), the trioma technique, the human B-cell
hybridoma technique (Kozbor *et al.*, Immunology Today (1983) 4:72) and
30 the EBV-hybridoma technique (Cole *et al.*, Monoclonal Antibodies and
Cancer Therapy, 77-96, Alan R. Liss, Inc., 1985).

Techniques for the production of single chain antibodies, such as those
described in U.S. Patent No. 4,946,778, can also be adapted to produce
35 single chain antibodies to polypeptides of this invention. Also, transgenic

- 15 -

mice, or other organisms, including other mammals, may be used to express humanized antibodies.

5 The above-described antibodies may be employed to isolate or to identify clones expressing the polypeptide or to purify the polypeptides by affinity chromatography. Antibodies against polypeptides of the present invention may also be employed to treat diseases of the invention, amongst others.

10 Polypeptides and polynucleotides of the present invention may also be used as vaccines. Accordingly, in a further aspect, the present invention relates to a method for inducing an immunological response in a mammal that comprises inoculating the mammal with a polypeptide of the present invention, adequate to produce antibody and/or T-cell immune response, including, for example, cytokine-producing T cells or cytotoxic T cells, to protect said animal from disease, whether that disease is already
15 established within the individual or not. An immunological response in a mammal may also be induced by a method comprises delivering a polypeptide of the present invention *via* a vector directing expression of the polynucleotide and coding for the polypeptide *in vivo* in order to induce such an immunological response to produce antibody to protect
20 said animal from diseases of the invention. One way of administering the vector is by accelerating it into the desired cells as a coating on particles or otherwise. Such nucleic acid vector may comprise DNA, RNA, a modified nucleic acid, or a DNA/RNA hybrid. For use a vaccine, a polypeptide or a nucleic acid vector will be normally provided as a vaccine formulation (composition). The formulation may further comprise
25 a suitable carrier. Since a polypeptide may be broken down in the stomach, it is preferably administered parenterally (for instance, subcutaneous, intramuscular, intravenous, or intradermal injection). Formulations suitable for parenteral administration include aqueous and
30 non-aqueous sterile injection solutions that may contain anti-oxidants, buffers, bacteriostats and solutes that render the formulation instonic with the blood of the recipient; and aqueous and non-aqueous sterile suspensions that may include suspending agents or thickening agents. The formulations may be presented in unit-dose or multi-dose containers,
35 for example, sealed ampoules and vials and may be stored in a freeze-dried condition requiring only the addition of the sterile liquid carrier

- 16 -

immediately prior to use. The vaccine formulation may also include adjuvant systems for enhancing the immunogenicity of the formulation, such as oil-in water systems and other systems known in the art. The dosage will depend on the specific activity of the vaccine and can be readily determined by routine experimentation.

Polypeptides of the present invention have one or more biological functions that are of relevance in one or more disease states, in particular the diseases of the invention hereinbefore mentioned. It is therefore useful to to identify compounds that stimulate or inhibit the function or level of the polypeptide. Accordingly, in a further aspect, the present invention provides for a method of screening compounds to identify those that stimulate or inhibit the function or level of the polypeptide. Such methods identify agonists or antagonists that may be employed for therapeutic and prophylactic purposes for such diseases of the invention as hereinbefore mentioned. Compounds may be identified from a variety of sources, for example, cells, cell-free preparations, chemical libraries, collections of chemical compounds, and natural product mixtures. Such agonists or antagonists so-identified may be natural or modified substrates, ligands, receptors, enzymes, etc., as the case may be, of the polypeptide; a structural or functional mimetic thereof (see Coligan *et al.*, Current Protocols in Immunology 1(2):Chapter 5 (1991)) or a small molecule.

The screening method may simply measure the binding of a candidate compound to the polypeptide, or to cells or membranes bearing the polypeptide, or a fusion protein thereof, by means of a label directly or indirectly associated with the candidate compound. Alternatively, the screening method may involve measuring or detecting (qualitatively or quantitatively) the competitive binding of a candidate compound to the polypeptide against a labeled competitor (e.g. agonist or antagonist). Further, these screening methods may test whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells bearing the polypeptide. Inhibitors of activation are generally assayed in the presence of a known agonist and the effect on activation by the agonist by the presence of the candidate compound is observed. Further, the screening methods may simply comprise the steps of mixing a candidate

- 17 -

compound with a solution containing a polypeptide of the present invention, to form a mixture, measuring a New GTPase-Activating Protein 1 activity in the mixture, and comparing the New GTPase-Activating Protein 1 activity of the mixture to a control mixture which contains no candidate compound.

Polypeptides of the present invention may be employed in conventional low capacity screening methods and also in high-throughput screening (HTS) formats. Such HTS formats include not only the well-established use of 96- and, more recently, 384-well microtiter plates but also emerging methods such as the nanowell method described by Schullek et al, Anal Biochem., 246, 20-29, (1997).

Fusion proteins, such as those made from Fc portion and New GTPase-Activating Protein 1 polypeptide, as hereinbefore described, can also be used for high-throughput screening assays to identify antagonists for the polypeptide of the present invention (see D. Bennett *et al.*, J Mol Recognition, 8:52-58 (1995); and K. Johanson *et al.*, J Biol Chem, 270(16):9459-9471 (1995)).

Screening techniques

The polynucleotides, polypeptides and antibodies to the polypeptide of the present invention may also be used to configure screening methods for detecting the effect of added compounds on the production of mRNA and polypeptide in cells. For example, an ELISA assay may be constructed for measuring secreted or cell associated levels of polypeptide using monoclonal and polyclonal antibodies by standard methods known in the art. This can be used to discover agents that may inhibit or enhance the production of polypeptide (also called antagonist or agonist, respectively) from suitably manipulated cells or tissues.

A polypeptide of the present invention may be used to identify membrane bound or soluble receptors, if any, through standard receptor binding techniques known in the art. These include, but are not limited to, ligand binding and crosslinking assays in which the polypeptide is labeled with a

radioactive isotope (for instance, ^{125}I), chemically modified (for instance, biotinylated), or fused to a peptide sequence suitable for detection or purification; and incubated with a source of the receptor (cells, cell membranes, cell supernatants, tissue extracts, bodily fluids). Other methods include biophysical techniques such as surface plasmon resonance and spectroscopy. These screening methods may also be used to identify agonists and antagonists of the polypeptide that compete with the binding of the polypeptide to its receptors, if any. Standard methods for conducting such assays are well understood in the art.

Examples of antagonists of polypeptides of the present invention include antibodies or, in some cases, oligonucleotides or proteins that are closely related to the ligands, substrates, receptors, enzymes, etc., as the case may be, of the polypeptide, e.g., a fragment of the ligands, substrates, receptors, enzymes, etc.; or a small molecule that bind to the polypeptide of the present invention but do not elicit a response, so that the activity of the polypeptide is prevented.

Screening methods may also involve the use of transgenic technology and New GTPase-Activating Protein 1 gene. The art of constructing transgenic animals is well established. For example, the New GTPase-Activating Protein 1 gene may be introduced through microinjection into the male pronucleus of fertilized oocytes, retroviral transfer into pre- or post-implantation embryos, or injection of genetically modified, such as by electroporation, embryonic stem cells into host blastocysts. Particularly useful transgenic animals are so-called "knock-in" animals in which an animal gene is replaced by the human equivalent within the genome of that animal. Knock-in transgenic animals are useful in the drug discovery process, for target validation, where the compound is specific for the human target. Other useful transgenic animals are so-called "knock-out" animals in which the expression of the animal ortholog of a polypeptide of the present invention and encoded by an endogenous DNA sequence in a cell is partially or completely annulled. The gene knock-out may be targeted to specific cells or tissues, may occur only in certain cells or tissues as a consequence of the limitations of the technology, or may occur in all, or substantially all, cells in the animal. Transgenic animal technology also offers a whole animal expression-cloning system in which introduced genes are expressed to give large amounts of polypeptides of the present invention

- 19 -

Screening kits for use in the above described methods form a further aspect of the present invention. Such screening kits compris :

- (a) a polypeptide of the present invention;
- (b) a recombinant cell expressing a polypeptide of the present invention;
- 5 (c) a cell membrane expressing a polypeptide of the present invention; or
- (d) an antibody to a polypeptide of the present invention;

which polypeptide is preferably that of SEQ ID NO:2.

It will be appreciated that in any such kit, (a), (b), (c) or (d) may comprise a substantial component.

10

Glossary

The following definitions are provided to facilitate understanding of certain terms used frequently hereinbefore.

15

"Antibodies" as used herein includes polyclonal and monoclonal antibodies, chimeric, single chain, and humanized antibodies, as well as Fab fragments, including the products of an

Fab or other immunoglobulin expression library.

20

"Isolated" means altered "by the hand of man" from its natural state, *i.e.*, if it occurs in nature, it has been changed or removed from its original environment, or both. For example, a polynucleotide or a polypeptide naturally present in a living organism is not "isolated," but the same polynucleotide or polypeptide separated from the coexisting materials of its natural state is "isolated", as the term is employed herein. Moreover, a polynucleotide or polypeptide that is introduced into an organism by transformation, genetic manipulation or by any other recombinant method

25

is "isolated" even if it is still present in said organism, which organism may be living or non-living.

"Polynucleotide" generally refers to any polyribonucleotide (RNA) or polydeoxribonucleotide (DNA), which may be unmodified or modified

RNA or DNA. "Polynucleotides" include, without limitation, single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or, more typically, double-stranded or a mixture of single- and double-stranded regions. In addition, "polynucleotide" refers to triple-stranded regions comprising RNA or DNA or both RNA and DNA. The term "polynucleotide" also includes DNAs or RNAs containing one or more modified bases and DNAs or RNAs with backbones modified for stability or for other reasons. "Modified" bases include, for example, tritylated bases and unusual bases such as inosine. A variety of modifications may be made to DNA and RNA; thus, "polynucleotide" embraces chemically, enzymatically or metabolically modified forms of polynucleotides as typically found in nature, as well as the chemical forms of DNA and RNA characteristic of viruses and cells. "Polynucleotide" also embraces relatively short polynucleotides, often referred to as oligonucleotides.

"Polypeptide" refers to any polypeptide comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids. "Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques that are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications may occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present to the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications. Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from post-translation natural processes or may be made by synthetic methods. Modifications include acetylation,

acylation, ADP-ribosylation, amidation, biotinylation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking, cyclization, disulfide bond formation, demethylation, formation of covalent cross-links, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination (see, for instance, *Proteins - Structure and Molecular Properties*, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993; Wold, F., *Post-translational Protein Modifications: Perspectives and Prospects*, 1-12, in *Post-translational Covalent Modification of Proteins*, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter *et al.*, "Analysis for protein modifications and nonprotein cofactors", *Meth Enzymol*, 182, 626-646, 1990, and Rattan *et al.*, "Protein Synthesis: Post-translational Modifications and Aging", *Ann NY Acad Sci*, 663, 48-62, 1992).

"Fragment" of a polypeptide sequence refers to a polypeptide sequence that is shorter than the reference sequence but that retains essentially the same biological function or activity as the reference polypeptide. "Fragment" of a polynucleotide sequence refers to a polynucleotide sequence that is shorter than the reference sequence of SEQ ID NO:1.

"Variant" refers to a polynucleotide or polypeptide that differs from a reference polynucleotide or polypeptide, but retains the essential properties thereof. A typical variant of a polynucleotide differs in nucleotide sequence from the reference polynucleotide. Changes in the nucleotide sequence of the variant may or may not alter the amino acid sequence of a polypeptide encoded by the reference polynucleotide. Nucleotide changes may result in amino acid substitutions, additions, deletions, fusions and truncations in the polypeptide encoded by the reference sequence, as discussed below. A typical variant of a polypeptide differs in amino acid sequence from the reference polypeptide. Generally, alterations are limited so that the sequences of the reference polypeptide and the variant are closely similar overall and, in many regions, identical. A variant and reference polypeptide may differ

in amino acid sequence by one or more substitutions, insertions, deletions in any combination. A substituted or inserted amino acid residue may or may not be one encoded by the genetic code. Typical conservative substitutions include Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe and Tyr. A variant of a polynucleotide or polypeptide may be naturally occurring such as an allele, or it may be a variant that is not known to occur naturally. Non-naturally occurring variants of polynucleotides and polypeptides may be made by mutagenesis techniques or by direct synthesis. Also included as variants are polypeptides having one or more post-translational modifications, for instance glycosylation, phosphorylation, methylation, ADP ribosylation and the like. Embodiments include methylation of the N-terminal amino acid, phosphorylations of serines and threonines and modification of C-terminal glycines.

"Allele" refers to one of two or more alternative forms of a gene occurring at a given locus in the genome.

"Polymorphism" refers to a variation in nucleotide sequence (and encoded polypeptide sequence, if relevant) at a given position in the genome within a population.

"Single Nucleotide Polymorphism" (SNP) refers to the occurrence of nucleotide variability at a single nucleotide position in the genome, within a population. An SNP may occur within a gene or within intergenic regions of the genome. SNPs can be assayed using Allele Specific Amplification (ASA). For the process at least 3 primers are required. A common primer is used in reverse complement to the polymorphism being assayed. This common primer can be between 50 and 1500 bps from the polymorphic base. The other two (or more) primers are identical to each other except that the final 3' base wobbles to match one of the two (or more) alleles that make up the polymorphism. Two (or more) PCR reactions are then conducted on sample DNA, each using the common primer and one of the Allele Specific Primers.

"Splice Variant" as used herein refers to cDNA molecules produced from RNA molecules initially transcribed from the same genomic DNA sequence but which have undergone alternative RNA splicing. Alternative RNA splicing occurs when a primary RNA transcript

und rgoes splicing, generally for the removal of introns, which results in the production of more than on mRNA molecule each of that may encode different amino acid sequences. The term splice variant also refers to the proteins encoded by the above cDNA molecules.

5 "Identity" reflects a relationship between two or more polypeptide sequences or two or more polynucleotide sequences, determined by comparing the sequences. In general, identity refers to an exact nucleotide to nucleotide or amino acid to amino acid correspondence of the two polynucleotide or two polypeptide sequences, respectively, over
10 the length of the sequences being compared.

"% Identity" - For sequences where there is not an exact correspondence, a "% identity" may be determined. In general, the two sequences to be compared are aligned to give a maximum correlation between the sequences. This may include inserting "gaps" in either one
15 or both sequences, to enhance the degree of alignment. A % identity may be determined over the whole length of each of the sequences being compared (so-called global alignment), that is particularly suitable for sequences of the same or very similar length, or over shorter, defined lengths (so-called local alignment), that is more suitable for sequences of
20 unequal length.

"Similarity" is a further, more sophisticated measure of the relationship between two polypeptide sequences. In general, "similarity" means a comparison between the amino acids of two polypeptide chains, on a residue by residue basis, taking into account not only exact
25 correspondences between a between pairs of residues, one from each of the sequences being compared (as for identity) but also, where there is not an exact correspondence, whether, on an evolutionary basis, one residue is a likely substitute for the other. This likelihood has an associated "score" from which the "% similarity" of the two sequences
30 can then be determined.

Methods for comparing the identity and similarity of two or more sequences are well known in the art. Thus for instance, programs available in the Wisconsin Sequence Analysis Package, version 9.1 (Devereux J et al, Nucleic Acids Res, 12, 387-395, 1984, available from
35 Genetics Computer Group, Madison, Wisconsin, USA), for example the

programs BESTFIT and GAP, may be used to determine the % identity between two polynucleotides and the % identity and the % similarity between two polypeptide sequences. BESTFIT uses the "local homology" algorithm of Smith and Waterman (J Mol Biol, 147,195-197, 1981, Advances in Applied Mathematics, 2, 482-489, 1981) and finds the best single region of similarity between two sequences. BESTFIT is more suited to comparing two polynucleotide or two polypeptide sequences that are dissimilar in length, the program assuming that the shorter sequence represents a portion of the longer. In comparison, GAP aligns two sequences, finding a "maximum similarity", according to the algorithm of Neddleman and Wunsch (J Mol Biol, 48, 443-453, 1970). GAP is more suited to comparing sequences that are approximately the same length and an alignment is expected over the entire length. Preferably, the parameters "Gap Weight" and "Length Weight" used in each program are 50 and 3, for polynucleotide sequences and 12 and 4 for polypeptide sequences, respectively. Preferably, % identities and similarities are determined when the two sequences being compared are optimally aligned.

Other programs for determining identity and/or similarity between sequences are also known in the art, for instance the BLAST family of programs (Altschul S F et al, J Mol Biol, 215, 403-410, 1990, Altschul S F et al, Nucleic Acids Res., 25:389-3402, 1997, available from the National Center for Biotechnology Information (NCBI), Bethesda, Maryland, USA and accessible through the home page of the NCBI at www.ncbi.nlm.nih.gov) and FASTA (Pearson W R, Methods in Enzymology, 183, 63-99, 1990; Pearson W R and Lipman D J, Proc Nat Acad Sci USA, 85, 2444-2448, 1988, available as part of the Wisconsin Sequence Analysis Package).

Preferably, the BLOSUM62 amino acid substitution matrix (Henikoff S and Henikoff J G, Proc. Nat. Acad Sci. USA, 89, 10915-10919, 1992) is used in polypeptide sequence comparisons including where nucleotide sequences are first translated into amino acid sequences before comparison.

Preferably, the program BESTFIT is used to determine the % identity of a query polynucleotide or a polypeptide sequence with respect to a reference polynucleotide or a polypeptide sequence, the query and the

reference sequence being optimally aligned and the parameters of the program set at the default value, as hereinbefore described.

"Identity Index" is a measure of sequence relatedness which may be used to compare a candidate sequence (polynucleotide or polypeptide) and a reference sequence. Thus, for instance, a candidate polynucleotide sequence having, for example, an Identity Index of 0.95 compared to a reference polynucleotide sequence is identical to the reference sequence except that the candidate polynucleotide sequence may include on average up to five differences per each 100 nucleotides of the reference sequence. Such differences are selected from the group consisting of at least one nucleotide deletion, substitution, including transition and transversion, or insertion. These differences may occur at the 5' or 3' terminal positions of the reference polynucleotide sequence or anywhere between these terminal positions, interspersed either individually among the nucleotides in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polynucleotide sequence having an Identity Index of 0.95 compared to a reference polynucleotide sequence, an average of up to 5 in every 100 of the nucleotides of the in the reference sequence may be deleted, substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

Similarly, for a polypeptide, a candidate polypeptide sequence having, for example, an Identity Index of 0.95 compared to a reference polypeptide sequence is identical to the reference sequence except that the polypeptide sequence may include an average of up to five differences per each 100 amino acids of the reference sequence. Such differences are selected from the group consisting of at least one amino acid deletion, substitution, including conservative and non-conservative substitution, or insertion. These differences may occur at the amino- or carboxy-terminal positions of the reference polypeptide sequence or anywhere between these terminal positions, interspersed either individually among the amino acids in the reference sequence or in one or more contiguous groups within the reference sequence. In other words, to obtain a polypeptide sequence having an Identity Index of 0.95 compared to a reference polypeptide sequence, an average of up to 5 in every 100 of the amino acids in the reference sequence may be deleted,

substituted or inserted, or any combination thereof, as hereinbefore described. The same applies *mutatis mutandis* for other values of the Identity Index, for instance 0.96, 0.97, 0.98 and 0.99.

5 The relationship between the number of nucleotide or amino acid differences and the Identity Index may be expressed in the following equation:

$$n_a \leq x_a - (x_a \bullet I),$$

in which:

n_a is the number of nucleotide or amino acid differences,

10 x_a is the total number of nucleotides or amino acids in SEQ ID NO:1 or SEQ ID NO:2, respectively,

I is the Identity Index ,

\bullet is the symbol for the multiplication operator, and

15 in which any non-integer product of x_a and I is rounded down to the nearest integer prior to subtracting it from x_a .

"Homolog" is a generic term used in the art to indicate a polynucleotide or polypeptide sequence possessing a high degree of sequence relatedness to a reference sequence. Such relatedness may be quantified by determining the degree of identity and/or similarity between the two
20 sequences as hereinbefore defined. Falling within this generic term are the terms "ortholog", and "paralog". "Ortholog" refers to a polynucleotide or polypeptide that is the functional equivalent of the polynucleotide or polypeptide in another species. "Paralog" refers to a polynucleotide or polypeptide that within the same species which is functionally similar.

25 "Fusion protein" refers to a protein encoded by two, unrelated, fused genes or fragments thereof. Examples have been disclosed in US 5541087, 5726044. In the case of Fc-NGAP1, employing an immunoglobulin Fc region as a part of a fusion protein is advantageous for performing the functional expression of Fc-NGAP1 or fragments of
30 NGAP1, to improve pharmacokinetic properties of such a fusion protein when used for therapy and to generate a dimeric NGAP1. The Fc-NGAP1

DNA construct comprises in 5' to 3' direction, a secretion cassette, i.e. a signal sequence that triggers export from a mammalian cell, DNA encoding an immunoglobulin Fc region fragment, as a fusion partner, and a DNA encoding NGAP1 or fragments thereof. In some uses it would be desirable to be able to alter the intrinsic functional properties (complement binding, Fc-Receptor binding) by mutating the functional Fc sides while leaving the rest of the fusion protein untouched or delete the Fc part completely after expression.

All publications and references, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference in their entirety as if each individual publication or reference were specifically and individually indicated to be incorporated by reference herein as being fully set forth. Any patent application to which this application claims priority is also incorporated by reference herein in its entirety in the manner described above for publications and references.

Claims

1. A polypeptide selected from the group consisting of:
 - (a) a polypeptide encoded by a polynucleotide comprising the sequence of SEQ ID NO:1;
 - 5 (b) a polypeptide comprising a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;
 - c) a polypeptide having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;
 - d) the polypeptide sequence of SEQ ID NO:2 and
 - 10 (e) fragments and variants of such polypeptides in (a) to (d).
2. The polypeptide of claim 1 comprising the polypeptide sequence of SEQ ID NO:2.
- 15 3. The polypeptide of claim 1 which is the polypeptide sequence of SEQ ID NO:2.
4. A polynucleotide selected from the group consisting of:
 - (a) a polynucleotide comprising a polynucleotide sequence having at least 95% identity to the polynucleotide sequence of SEQ ID NO:1;
 - 20 (b) a polynucleotide having at least 95% identity to the polynucleotide of SEQ ID NO:1;
 - (c) a polynucleotide comprising a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;
 - 25 (d) a polynucleotide having a polynucleotide sequence encoding a polypeptide sequence having at least 95% identity to the polypeptide sequence of SEQ ID NO:2;

- (e) a polynucleotide with a nucleotide sequence of at least 100 nucleotides obtained by screening a library under stringent hybridization conditions with a labeled probe having the sequence of SEQ ID NO: 1 or a fragment thereof having at least 15 nucleotides;
- 5 (f) a polynucleotide which is the RNA equivalent of a polynucleotide of (a) to (e);
- (g) a polynucleotide sequence complementary to said polynucleotide of any one of (a) to (f), and
- (h) polynucleotides that are variants or fragments of the polynucleotides of any one of (a) to (g) or that are complementary to above mentioned polynucleotides,
- 10 over the entire length thereof.

5. A polynucleotide of claim 4 selected from the group consisting of:

- (a) a polynucleotide comprising the polynucleotide of SEQ ID NO:1;
- (b) the polynucleotide of SEQ ID NO:1;
- 15 (c) a polynucleotide comprising a polynucleotide sequence encoding the polypeptide of SEQ ID NO:2; and
- (d) a polynucleotide encoding the polypeptide of SEQ ID NO:2.

6. An expression system comprising a polynucleotide capable of producing a polypeptide of any one of claim 1-3 when said expression vector is present in a compatible host cell.

20

7. A recombinant host cell comprising the expression vector of claim 6 or a membrane thereof expressing the polypeptide of any one of claim 1-3.

25

8. A process for producing a polypeptide of any one of claim 1-3 comprising the step of culturing a host cell as defined in claim 7 under conditions sufficient for

th production of said polypeptide and recovering the polypeptide from the culture medium.

9. A fusion protein consisting of the Immunoglobulin Fc-region and a polypeptide
5 any one of claims 1-3.

10. An antibody immunospecific for the polypeptide of any one of claims 1 to 3.

11. A method for screening to identify compounds that stimulate or inhibit the function or level of the polypeptide of any one of claim 1-3 comprising a method selected from the group consisting of:

10 (a) measuring or, detecting, quantitatively or qualitatively, the binding of a candidate compound to the polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof by means of a label directly or indirectly associated with the candidate compound;

(b) measuring the competition of binding of a candidate compound to the
15 polypeptide (or to the cells or membranes expressing the polypeptide) or a fusion protein thereof in the presence of a labeled competitor;

(c) testing whether the candidate compound results in a signal generated by activation or inhibition of the polypeptide, using detection systems appropriate to the cells or cell membranes expressing the polypeptide;

20 (d) mixing a candidate compound with a solution containing a polypeptide of any one of claims 1-3, to form a mixture, measuring activity of the polypeptide in the mixture, and comparing the activity of the mixture to a control mixture which contains no candidate compound; or

(e) detecting the effect of a candidate compound on the production of mRNA
25 encoding said polypeptide or said polypeptide in cells, using for instance, an ELISA assay, and

(f) producing said compound according to biotechnological or chemical standard techniques.

- 1 -

SEQUENCE LISTING

<110> Merck Patent GmbH

5 <120> New GTPase activating protein 1

<130> NGAP1CSWS

<140>

10 <141>

<160> 2

<170> PatentIn Ver. 2.1

15

<210> 1

<211> 1836

<212> DNA

<213> Homo sapiens

20

 $\langle 220 \rangle$

<221> CDS

<222> (80) .. (1486)

25 <400> 1

ttgaggggtgt gactccccag gaccagggag atgtttatgt ccaactgata acatatatttta 60

ctctcttctc caggatgac atg gaa gct tac cgg acc caq aac tgc ttc ctc 112

30	Met	Glu	Ala	Tyr	Arg	Thr	Gln	Asn	Cys	Phe	Leu
	1				5					10	

aac tcc gag atc cac cag gtc aca aag atc tgg aga aaq gtg gct gaq 160

Asn Ser Glu Ile His Gln Val Thr Lys Ile Trp Arg Lys Val Ala Glu
15 20 25

35

aag gag aag gcc ctt ctg acg aag tgc gcc tac ctc caa gcc aga aac 208

Lys Glu Lys Ala Leu Leu Thr Lys Cys Ala Tyr Leu Gln Ala Arg Asn
30 35 40

40 tgc cag gtg gaa agc aag tac ctg gcc ggt ctg aga agg ctg cag gac 256

Cys Gln Val Glu Ser Lys Tyr Leu Ala Gly Leu Arg Arg Leu Gln Glu

gcc ctg ggg gac gaa gcc agc gag tgc tca gag ctg ctg agg cag ctt 304

Ala Leu Gly Asp Glu Ala Ser Glu Cys Ser Glu Leu Leu Arg Gln Leu
60 65 70 75

gtc cag gag gca ctg cag tgg gaa gct ggg gag gcc tca tct gac agc 352

Val Gln Glu Ala Leu Gln Trp Glu Ala Gly Glu Ala Ser Ser Asp Ser
80 85 90

atc gag ctg agc ccc atc agt aag tat gat gag tac ggc ttc ctg acg 400

Ile Glu Leu Ser Pro Ile Ser Lys Tyr Asp Glu Tyr Gly Phe Leu Thr
95 100 105

55

gtg ccc gac tat gag gtg gaa gac ctg aag ctg ctg gcc aag atc caa 448

Val	Pro	Asp	Tyr	Glu	Val	Glu	Asp	Leu	Lys	Leu	Leu	Ala	Lys	Ile	Gln
		110					115					120			

60

- 2 -

	gca ttg gag tca cga tcc cac cac ctg ctg ggc ctc gag gct gtg gat	496
	Ala Leu Glu Ser Arg Ser His His Leu Leu Gly Leu Glu Ala Val Asp	
	125 130 135	
5	cgg ccg ctg agg gag cgc tgg gct gcc ctg ggc gat ctt gtg ccc tca	544
	Arg Pro Leu Arg Glu Arg Trp Ala Ala Leu Gly Asp Leu Val Pro Ser	
	140 145 150 155	
10	gcc gag ctc aag cag cta ctg cgg gca gga gta ccc cgt gaa cac cgg	592
	Ala Glu Leu Lys Gln Leu Leu Arg Ala Gly Val Pro Arg Glu His Arg	
	160 165 170	
15	cct cgt gtc tgg agg tgg ctg gtc cac ctc cgt gtc cag cac ctg cac	640
	Pro Arg Val Trp Arg Trp Leu Val His Leu Arg Val Gln His Leu His	
	175 180 185	
20	act cca ggc tgc tac cag gaa ctg ctg agc cgg ggc cag gcc cgc gag	688
	Thr Pro Gly Cys Tyr Gln Glu Leu Leu Ser Arg Gly Gln Ala Arg Glu	
	190 195 200	
	cac cct gct gcc cgc cag att gag ctg gac ctg aac cgg acc ttc ccc	736
	His Pro Ala Ala Arg Gln Ile Glu Leu Asp Leu Asn Arg Thr Phe Pro	
	205 210 215	
25	aac aac aaa cac ttc acc tgc ccc acc tcc agc ttc ccc gac aag ctc	784
	Asn Asn Lys His Phe Thr Cys Pro Thr Ser Ser Phe Pro Asp Lys Leu	
	220 225 230 235	
30	cgc cgg gtg ctg ctg gcc ttc tcc tgg cag aac ccc acc atc ggc tac	832
	Arg Arg Val Leu Leu Ala Phe Ser Trp Gln Asn Pro Thr Ile Gly Tyr	
	240 245 250	
35	tgc cag ggc ctg aac agg ctg gcg gcc att gcc ctg ctg gtc cta gag	880
	Cys Gln Gly Leu Asn Arg Leu Ala Ala Ile Ala Leu Leu Val Leu Glu	
	255 260 265	
40	gag gag gag agc gcc ttc tgg tgc ctg gtg gcc att gtg gag acc atc	928
	Glu Glu Glu Ser Ala Phe Trp Cys Leu Val Ala Ile Val Glu Thr Ile	
	270 275 280	
	atg ccc gct gat tac tac tgc aac acg ctg acg gca tcc cag gtg gac	976
	Met Pro Ala Asp Tyr Tyr Cys Asn Thr Leu Thr Ala Ser Gln Val Asp	
	285 290 295	
45	cag cgg gtg ctc cag gac ctg ctc tcg gag aag ctg ccc agg ctg atg	1024
	Gln Arg Val Leu Gln Asp Leu Leu Ser Glu Lys Leu Pro Arg Leu Met	
	300 305 310 315	
50	gcc cat ctg ggg cag cac cac gtg gat ctc tcc ctc gtc acc ttc aac	1072
	Ala His Leu Gly Gln His His Val Asp Leu Ser Leu Val Thr Phe Asn	
	320 325 330	
55	tgg ttc ctc gtg gtc ttt gcg gac agt ctc att agc aac atc ctc ctt	1120
	Trp Phe Leu Val Val Phe Ala Asp Ser Leu Ile Ser Asn Ile Leu Leu	
	335 340 345	
60	cgg gtc tgg gat gcc ttc ctg tac gag ggg acg aag gtg gtg ttt cgc	1168
	Arg Val Trp Asp Ala Phe Leu Tyr Glu Gly Thr Lys Val Val Phe Arg	
	350 355 360	

- 3 -

tat gcc ttg gcc att ttc aag tac aac gag aag gag atc ttg agg cta 1216
 Tyr Ala Leu Ala Ile Phe Lys Tyr Asn Glu Lys Glu Ile Leu Arg L u
 365 370 375

5 cag aat ggc ctg gaa atc tac cag tac ctg cgc ttc ttc acc aag acc 1264
 Gln Asn Gly Leu Glu Ile Tyr Gln Tyr Leu Arg Phe Phe Thr Lys Thr
 380 385 390 395

10 atc tcc aac agc cgg aag ctg atg aac atc gcc ttc aat gac atg aac 1312
 Ile Ser Asn Ser Arg Lys Leu Met Asn Ile Ala Phe Asn Asp Met Asn
 400 405 410

15 ccc ttc cgc atg aaa cag ctg cgg cag ctg cgc atg gtc cac cgg gag 1360
 Pro Phe Arg Met Lys Gln Leu Arg Gln Leu Arg Met Val His Arg Glu
 415 420 425

20 cgg ctg gag gct gag ctg cgg gag ctg gag cag ctt aag gca gag tac 1408
 Arg Leu Glu Ala Glu Leu Arg Glu Leu Glu Gln Leu Lys Ala Glu Tyr
 430 435 440

25 ctg gag agg cgg gca tcc cgg cgc aga gct gtg tcc gag ggc tgt gcc 1456
 Leu Glu Arg Arg Ala Ser Arg Arg Arg Ala Val Ser Glu Gly Cys Ala
 445 450 455

30 agc gag gac gag gtg gag ggg gaa gcc tga cttggccacc tcccctcccc 1506
 Ser Glu Asp Glu Val Glu Gly Glu Ala
 460 465

35 acagccttcc tcacccttgg ctggcagacc cactggaggt caggcacgga ccagtggccc 1566
 agccctgggt gtcccatcac catgtgacct tggacatgtc ccttcccctc tctggccctc 1626
 agtttcccca ctgggacatt gtgtgctgca aagccattgg ttgggctact tcttcatagg 1686

40 cacttactta ccagggatg ccacccttcc gtcacctctt ccacagagca ctttggcatg 1746
 taaacaagca agagcactgc ctctataggg taacctggaa cattcttttag gttatatcaa 1806
 tataaaacaa tgtaaattgt ggaaatcatt 1836

45 <210> 2
 <211> 468
 <212> PRT
 <213> Homo sapiens

50 <400> 2
 Met Glu Ala Tyr Arg Thr Gln Asn Cys Phe Leu Asn Ser Glu Ile His
 1 5 10 15
 Gln Val Thr Lys Ile Trp Arg Lys Val Ala Glu Lys Glu Lys Ala Leu
 20 25 30
 Leu Thr Lys Cys Ala Tyr Leu Gln Ala Arg Asn Cys Gln Val Glu Ser
 35 40 45
 Lys Tyr Leu Ala Gly Leu Arg Arg Leu Gln Glu Ala Leu Gly Asp Glu
 50 55 60
 Ala Ser Glu Cys Ser Glu Leu Leu Arg Gln Leu Val Gln Glu Ala Leu
 65 70 75 80
 Gln Trp Glu Ala Gly Glu Ala Ser Ser Asp Ser Ile Glu Leu Ser Pro
 85 90 95
 60 Ile Ser Lys Tyr Asp Glu Tyr Gly Phe Leu Thr Val Pro Asp Tyr Glu
 100 105 110

	Val	Glu	Asp	Leu	Lys	Leu	Leu	Ala	Lys	Ile	Gln	Ala	Leu	Glu	Ser	Arg
			115					120					125			
	Ser	His	His	Leu	Leu	Gly	Leu	Glu	Ala	Val	Asp	Arg	Pro	Leu	Arg	Glu
		130					135					140				
5	Arg	Trp	Ala	Ala	Leu	Gly	Asp	Leu	Val	Pro	Ser	Ala	Glu	Leu	Lys	Gln
	145					150					155					160
	Leu	Leu	Arg	Ala	Gly	Val	Pro	Arg	Glu	His	Arg	Pro	Arg	Val	Trp	Arg
					165					170					175	
10	Trp	Leu	Val	His	Leu	Arg	Val	Gln	His	Leu	His	Thr	Pro	Gly	Cys	Tyr
				180					185					190		
	Gln	Glu	Leu	Leu	Ser	Arg	Gly	Gln	Ala	Arg	Glu	His	Pro	Ala	Ala	Arg
			195					200					205			
	Gln	Ile	Glu	Leu	Asp	Leu	Asn	Arg	Thr	Phe	Pro	Asn	Asn	Lys	His	Phe
		210					215					220				
15	Thr	Cys	Pro	Thr	Ser	Ser	Phe	Pro	Asp	Lys	Leu	Arg	Arg	Val	Leu	Leu
	225					230					235					240
	Ala	Phe	Ser	Trp	Gln	Asn	Pro	Thr	Ile	Gly	Tyr	Cys	Gln	Gly	Leu	Asn
					245					250					255	
20	Arg	Leu	Ala	Ala	Ile	Ala	Leu	Leu	Val	Leu	Glu	Glu	Glu	Glu	Ser	Ala
				260					265					270		
	Phe	Trp	Cys	Leu	Val	Ala	Ile	Val	Glu	Thr	Ile	Met	Pro	Ala	Asp	Tyr
			275					280					285			
	Tyr	Cys	Asn	Thr	Leu	Thr	Ala	Ser	Gln	Val	Asp	Gln	Arg	Val	Leu	Gln
		290					295					300				
25	Asp	Leu	Leu	Ser	Glu	Lys	Leu	Pro	Arg	Leu	Met	Ala	His	Leu	Gly	Gln
	305					310					315					320
	His	His	Val	Asp	Leu	Ser	Leu	Val	Thr	Phe	Asn	Trp	Phe	Leu	Val	Val
				325						330					335	
30	Phe	Ala	Asp	Ser	Leu	Ile	Ser	Asn	Ile	Leu	Leu	Arg	Val	Trp	Asp	Ala
				340				345						350		
	Phe	Leu	Tyr	Glu	Gly	Thr	Lys	Val	Val	Phe	Arg	Tyr	Ala	Leu	Ala	Ile
			355					360					365			
	Phe	Lys	Tyr	Asn	Glu	Lys	Glu	Ile	Leu	Arg	Leu	Gln	Asn	Gly	Leu	Glu
		370					375					380				
35	Ile	Tyr	Gln	Tyr	Leu	Arg	Phe	Phe	Thr	Lys	Thr	Ile	Ser	Asn	Ser	Arg
	385					390					395					400
	Lys	Leu	Met	Asn	Ile	Ala	Phe	Asn	Asp	Met	Asn	Pro	Phe	Arg	Met	Lys
				405						410					415	
40	Gln	Leu	Arg	Gln	Leu	Arg	Met	Val	His	Arg	Glu	Arg	Leu	Glu	Ala	Glu
				420					425					430		
	Leu	Arg	Glu	Leu	Glu	Gln	Leu	Lys	Ala	Glu	Tyr	Leu	Glu	Arg		

INTERNATIONAL SEARCH REPORT

Intern: Application No

PCT/EP 01/04888

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/12 C07K14/47 C12N15/62 C07K16/18 G01N33/68
C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K G01N C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE EM_HUM 'Online! EMBL; Accession Number: AL353935, 27 April 2000 (2000-04-27) BLOECKER H. ET. AL.: XP002176109 the whole document	1-11
P,X	DATABASE SWALL 'Online! Accession NO: Q9NSH4, 1 October 2000 (2000-10-01) BLOECKER H. ET AL: XP002176110 the whole document	1-11
	--- -/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *8* document member of the same patent family

Date of the actual completion of the international search

29 August 2001

Date of mailing of the international search report

10/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Madruga, J

INTERNATIONAL SEARCH REPORT

Intern Application No

PCT/EP 01/04888

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	SCHEFFZEK K ET AL: "GTPase-activating proteins: helping hands to complement an active site" TIBS TRENDS IN BIOCHEMICAL SCIENCES, ELSEVIER PUBLICATION, CAMBRIDGE, EN, vol. 23, no. 7, 1 July 1998 (1998-07-01), pages 257-262, XP004126978 ISSN: 0968-0004 the whole document	1-11
A	NEUWALD A F: "A shared domain between a spindle assembly checkpoint protein and Ypt/Rab-specific GTPase-activators" TIBS TRENDS IN BIOCHEMICAL SCIENCES, ELSEVIER PUBLICATION, CAMBRIDGE, EN, vol. 22, no. 7, 1 July 1997 (1997-07-01), pages 243-244, XP004081590 ISSN: 0968-0004 cited in the application the whole document	1-11
A	EP 0 496 162 A (MERCK & CO INC) 29 July 1992 (1992-07-29) page 3, line 15 - line 41; examples VII, VIII	11

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern: Application No

PCT/EP 01/04888

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0496162 A	29-07-1992	CA 2057675 A JP 5178894 A	25-06-1992 20-07-1993